

Comparative Study of Mental Stress Score between Various Surgical Modalities for Treatment of 10–20 mm Renal Calculi

Velmurugan Palaniyandi¹, Siddharth Munisamy², Sriram Krishnamoorthy³, Venkat Ramanan³, Natarajan Kumaresan³

¹Assistant Professor, Department of Urology, Sri Ramachandra Medical College and Research Institute, Chennai, Tamil Nadu, India,

²Consultant Urologist, KDR Hospital, Thanjavur, Tamil Nadu, India, ³Professor of Urology, Department of Urology, Sri Ramachandra Medical College and Research Institute, Chennai, Tamil Nadu, India

Abstract

Introduction: The treatment of renal calculi has undergone a paradigm change in the past two decades. The ultimate aim in treating these stones will be to achieve a complete clearance with minimal morbidity. The risk of recurrence or incomplete clearance or need for ancillary procedures would significantly affect patients' mental health.

Purpose: The purpose of this study is to compare mental stress experienced by the patient on the completion of the treatment using Kessler Psychological Distress Scale (K10) for each modality in the management of renal and upper ureteric calculi 10–20 mm size.

Materials and Methods: This was a prospective study on patients with renal calculi. Following the procedure, patients were asked to fill up and complete the questionnaire at 2 and 6 weeks. The K10 questionnaire contained 10 questions with each of them having 5 outcomes. Based on the total score, the degrees of mental distress due to the procedure were graded as mild, moderate, and severe.

Results: When assessing patients' psychological status using Kessler's scale at 2 weeks, a majority of patients had a score of <20. However, one-fifth of patients in the retrograde intrarenal surgery group and about one-sixth of patients in percutaneous nephrolithotomy (PCNL) group had a mild distress score. When assessing patients' psychological status using Kessler's scale at 6 weeks following primary procedure, most patients are <20 score, and hence, not statistically significant. But when the Shock wave lithotripsy (SWL) group was analyzed, 2.5% of patients had a significant psychological stress.

Conclusion: All three procedures have a good acceptance rate among the patients. PCNL, though more invasive than the other two procedures, has a better acceptance rate in view of achieving complete clearance in one sitting. On the other hand, extracorporeal SWL (ESWL), despite being so non-invasive, has a significant psychological impact in view of the need for multiple sittings and associated complications.

Key words: Lithotripsy, nephrolithotomy, questionnaire, renal calculi

INTRODUCTION

The primary goal while treating renal calculi and upper ureteric calculi is to achieve maximum clearance of stone with minimal morbidity. The various minimally invasive modalities described for treatment of such

stones are shockwave lithotripsy (SWL), percutaneous nephrolithotomy (PCNL), and retrograde intrarenal surgery (RIRS).^{1,2} The same authors had earlier published their study comparing success rate, retreatment rate, need for auxiliary procedure, complication rate, mean procedure time, and mean hospital stay between the above three modalities. As a part of that study, we also had assessed the mental stress score following the surgical treatment received using Kessler Psychological Distress Scale (K10). Kessler Psychological Distress Scale (K10) is used to assess the patients' mental stress, which was proposed by Kessler R, Professor of health care policy, Harvard Medical School, Boston, USA. This K 10 test is a patient's self-report measure to assess their current condition about anxiety and depression following

Access this article online



www.ijss-sn.com

Month of Submission : 12-2017

Month of Peer Review : 01-2018

Month of Acceptance : 01-2018

Month of Publishing : 02-2018

Corresponding Author: Dr. Sriram Krishnamoorthy, Department of Urology, Sri Ramachandra Medical College, Chennai – 600 116, Tamil Nadu, India. Phone: +91-8056139257. E-mail: sriramuro@gmail.com

a procedure.^[3] The 10-item Kessler Psychological Distress Scale (K10) is a short measure of non-specific psychological distress, which has been shown to be a sensitive screen for the Diagnostic and Statistical Manual of Mental Disorders criteria for anxiety and mood disorders.^[4]

The three surgical procedures taken for consideration in our study are all very well established and widely accepted modalities of treatment. For a stone smaller than 10 mm, initial treatment option has always been extracorporeal SWL. For a large stone above 20 mm, PCNL is the accepted modality of treatment. For stones in the interim range, the options are plenty. The level of acceptance of these procedures also varies from one procedure to the other. The purpose of this study is to identify if any of the above-mentioned procedures have any impact on the ultimate psychological status that might alter the final outcome in such patients.

Aim

The aim of this study is to compare mental stress experienced by the patient on the completion of the treatment using Kessler Psychological Distress Scale (K10) for each modality in the management of renal and upper ureteric calculi 10–20 mm size.

MATERIALS AND METHODS

This was a prospective observational comparative study conducted in a teaching institution over 1 year period from November 2013 to October 2014. A total of 287 cases renal and upper ureteric calculi of 10–20 mm size were included in our study. All patients underwent PCNL, RIRS, or SWL. The study model was presented to the Institutional Ethical Committee and approval was obtained. Following the procedure, patients were asked to fill up and complete the questionnaire at 2 and 6 weeks. The K10 questionnaire contained 10 questions with each of them having 5 outcomes. The outcomes were numbered and rated from 1 to 5.

Table 1 describes the details of the K 10 questionnaire circulated among the patients at 2 and 6 weeks after

treatment. Based on the total score, the degrees of mental distress due to the procedure were graded as follows:

Score <20, where patients are likely to be well without any mental illness.

Score 20–24, where patients are likely to have mild mental disorder.

Score 25–29, where patients are likely to have moderate mental disorder.

Score >29, where patients are likely to have serious mental disorder.

Inclusion and Exclusion Criteria

All patients presenting with calculi of 10–20 mm in otherwise normal renal pelvicalyceal system, pelviureteric junction, or proximal ureter up to L3 transverse process were included in our study. Those patients in whom the stone was below L3 transverse process, multiple renal calculi with second calculi size more than 4 mm, abnormal upper urinary tract anatomy such as duplex system, horseshoe kidney, ectopic kidney, and pelviureteric junction obstruction, and patients with axial skeletal abnormality such as scoliosis and kyphosis or bleeding diathesis were all excluded from the study.

The choice of treatment modality for the management of the upper urinary tract calculi of 10–20 mm calculi is largely determined by the individual surgeon taking into consideration the patient’s anatomy, comorbid conditions, urinary tract anatomy, stone density, and location as well as patients’ preference.

Statistical Analysis

Descriptive statistical analysis was carried out in the present study. To describe about the data frequency analysis, percentage analysis was used for categorical variables, and for continuous variables, the mean and standard deviation were used. For the multivariate analysis, the Kruskal–Wallis test and ANOVA were used, and for trivariate and bivariate analysis, Mann–Whitney test was used. To find the significance in categorical data, Chi-square test was used. In both the above statistical tools, the probability value of <0.05 is considered as significant level. The statistical

Table 1: K 10 test questionnaire given to patients at 2 and 6 weeks after treatment

Questionnaire	Scoring points				
During the past 15 days, about how often did you feel tires out of no good reason?	1	2	3	4	5
During the past 15 days, about how often did you feel nervous?	1	2	3	4	5
During the past 15 days, about how often did you feel so nervous that nothing could calm you down?	1	2	3	4	5
During the past 15 days, about how often did you feel hopeless?	1	2	3	4	5
During the past 15 days, about how often did you feel restless and fidgety?	1	2	3	4	5
During the past 15 days, about how often did you feel so restless you could not sit still?	1	2	3	4	5
During the past 15 days, about how often did you feel depressed?	1	2	3	4	5
During the past 15 days, about how often did you feel everything was an effort?	1	2	3	4	5
During the past 15 days, about how often did you feel so sad that nothing could cheer you up?	1	2	3	4	5
During the past 15 days, about how often did you feel worthless?	1	2	3	4	5

1 - None of the time, 2 - A little of the time, 3 - Some of the time, 4 - Most of the time, 5 - All the time

software SPSS 16.0 version was used for the analysis of the data, and Microsoft Word and Excel have been used to generate graphs and tables.

Observations

Table 2 and Figure 1 illustrates the Kessler’s distress scale at 2 weeks after intervention. When assessing patients’ psychological status using Kessler’s scale at 2 weeks, a majority of patients had a score of <20. However, one-fifth of patients in the RIRS group and about one-sixth of patients in PCNL group had a mild distress score. None of the patients in RIRS group had moderate or severe distress scores.

Table 3 and Figure 2 illustrates the Kessler’s distress scale at 6 weeks after intervention. When assessing patients’ psychological status using Kessler’s scale at 6 weeks following primary procedure, most patients are <20 score and hence not statistical significant. But when the Shock wave lithotripsy (SWL) group was analyzed, 2.5% of patients had a significant psychological stress.. Following PCNL as the primary treatment at the end of the study at 6 weeks, all patients are stone free and without mental distress. In RIRS group, the mental distress at the end of 6 weeks is within the comfort zone. In SWL group, the mental distress at the end of 6 weeks is within the comfort zone in almost all patients with 2 patients having mild distress at 6 weeks in view of prolonged need of completion of stone management.

DISCUSSION

Renal calculus and the bothersome symptoms increase the need for multiple visits to hospitals and also cause significant psychological impact on patients.^[5] Moreover, continuous usage of medication, need for recurrent

surgical procedures, limitation in physical activity, and loss of working hours and associated financial burden are found to be associated with many psychological disturbances. Among these, anxiety has been identified as the more prevalent entity and is associated with poor patient compliance toward management and even progress to depression in many patients.^[6,7]

In the treatment of renal stones, the type of anesthesia, need for an inpatient admission, success and failure rate of the procedure, complications, need for the auxiliary procedure, and affordability to the procedures all play a role in influencing patients’ anxiety status.




Figure 1: Kessler psychological distress scale at 2 weeks




Figure 2: Kessler psychological distress scale at 6 weeks

Table 2: Kessler psychological distress scale at 2 weeks

Distress score	Group, n (%)				Chi-square	P value
	PCNL	RIRS	ESWL	Total		
Below 20	132 (82)	36 (80)	67 (82.7)	235 (81.9)	7.535	0.110
20–24	22 (13.7)	9 (20)	7 (8.6)	38 (13.2)		
25–29	7 (4.3)	0	7 (8.6)	14 (4.9)		
30 and above	0	0	0	0		

PCNL: Percutaneous nephrolithotomy, RIRS: Retrograde intrarenal surgery, ESWL: Extracorporeal shockwave lithotripsy

Table 3: Kessler psychological distress scale at 6 weeks

Distress score	Group, n (%)				Chi-square	P value
	PCNL	RIRS	ESWL	Total		
Below 20	161 (100)	45 (100)	79 (97.5)	285 (99.3)	5.122	0.077
20–24	0	0	2 (2.5)	2 (0.7)		
25–29	0	0	0	0		
30 and above	0	0	0	0		

PCNL: Percutaneous nephrolithotomy, RIRS: Retrograde intrarenal surgery, ESWL: Extracorporeal shockwave lithotripsy

In our earlier study when assessing the failure rates of these procedures, SWL recorded the maximum failure (17.3%), with 19.8% of patients needing auxiliary procedure.^[8] On the other hand, PCNL and RIRS (5.6% and 11.1%, respectively) had the least number of cases needing an auxiliary procedure.^[8]

The degrees of mental distress due to the procedure were graded based on Andrew's study.^[9] The work done by Brown *et al.* assessed patients' anxiety before and after different surgical procedures for renal calculus and to identify various factors that can attribute to pre- and post-operative anxiousness.^[10] The measurement of palmar sweat (evaporimeter) for quantifying stress and bipolar visual analog scale and the Spielberger State Anxiety Questionnaire were used before and after the procedures. The study concluded highly significant reduction in the palmar sweat production and score obtained following open surgery, but no changes in patients undergoing PCNL or lithotripsy before and after treatment. Pre-operatively, fear of a general anesthetic, and post-operatively, pain were identified as cause for patients' anxiety.

State and Trait Anxiety Inventory scale was used to assess the anxiety status of 128 patients following SWL. It showed that the residual fragments after SWL procedure made patients more anxious as additional procedures were needed for clearing the left out stones. The study also concluded that detailed information should be provided to patients with respect to procedure and possible complications, and potential need for additional treatment may reduce the anxiety.^[11]

A study done by Brown *et al.* with 24 patients undergoing extracorporeal piezolithotripsy with continuous assessment of anxiety by measurement of palmar sweat during the procedure itself found with 50% of patients showed increased levels of palmar sweat throughout treatment, with a return to pre-treatment levels after the procedure. 8 patients attributed pain as their cause for anxiety. Patient education before the procedure may help in reducing anxiety.^[12]

Extracorporeal SWL (ESWL) seems to be safe and simple and being done as an outpatient procedure. It does not require any anesthesia and patient can go back home the same day. All of these factors support the fact that it will not have any impact on the psychological status of the patients, but it was not so in our study. This study underlines the fact that the patient anxiety depends not only on the severity of the procedure but also on the sequelae of the procedure, such as incomplete clearance, complications of the procedure, need for multiple admissions, and ancillary procedures.

Limitations of this Study

The sample size is small and associated with difficulty in performing stratified analysis. The study was unable to exclude many of the other confounding factors which may have influenced some of the outcomes analyzed which is beyond the scope and purview of this study.

CONCLUSION

All three procedures have a good acceptance rate among the patients. PCNL, though more invasive than the other two procedures, has a better acceptance rate in view of achieving complete clearance in one sitting. On the other hand, ESWL, despite being so non-invasive, has a significant psychological impact in view of the need for multiple sittings and associated complications.

REFERENCES

1. Srisubath A, Potisat S, Lojanapiwat B, Setthawong V, Laopaiboon M. Extracorporeal shock wave lithotripsy (ESWL) versus percutaneous nephrolithotomy (PCNL) or retrograde intrarenal surgery (RIRS) for kidney stones. *Cochrane Database Syst Rev* 2009;4:CD007044.
2. Turk C, Knoll T, Petrik A, Sarica K, Straub M, Seitz C. Guidelines on Urolithiasis. *European Association of Urology Update*; 2011.
3. Kessler RC, Andrews G, Colpe LJ, Hiripi E, Mroczek DK, Normand SL, *et al.* Short screening scales to monitor population prevalences and trends in non-specific psychological distress. *Psychol Med* 2002;32:959-76.
4. Bougie E, Arim RG, Kohen DE, Findlay LC. Validation of the 10-item kessler psychological distress scale (K10) in the 2012 aboriginal peoples survey. *Health Rep* 2016;27:3-10.
5. Herrman H, Patrick DL, Diehr P, Martin ML, Fleck M, Simon GE, *et al.* Longitudinal investigation of depression outcomes in primary care in six countries: The LIDO study. Functional status, health service use and treatment of people with depressive symptoms. *Psychol Med* 2002;32:889-902.
6. Greenberg PE, Leong SA, Birnbaum HG, Robinson RL. The economic burden of depression with painful symptoms. *J Clin Psychiatry* 2003;64 Suppl 7:17-23.
7. Alonso J, Ferrer M, Gandek B, Ware JE Jr, Aaronson NK, Mosconi P, *et al.* Health-related quality of life associated with chronic conditions in eight countries: Results from the international quality of life assessment (IQOLA) project. *Qual Life Res* 2004;13:283-98.
8. Muniasamy S, Palaniyandi V, Krishnamoorthy S, Kumaresan N, Ramanan V. Comparative study of stone free rate, morbidity and need for retreatment procedures between various surgical modalities for 10-20 mm upper urinary tract calculi. *Int J Sci Study* 2017;4:37-43.
9. Andrews G, Slade T. Interpreting scores on the kessler psychological distress scale (K10). *Aust N Z J Public Health* 2001;25:494-7.
10. Brown SM. Quantitative measurement of anxiety in patients undergoing surgery for renal calculus disease. *J Adv Nurs* 1990;15:962-70.
11. Şahin C, Sarica K. Comparative analysis of shock wave therapy success rate in management of renal stones and patient anxiety. *South Clin Ist Euras*. 2017;28:47-52.
12. Brown SM. Peri-operative anxiety in patients undergoing extracorporeal piezolithotripsy. *J Adv Nurs* 1990;15:1078-82.

How to cite this article: Palaniyandi V, Muniasamy S, Krishnamoorthy S, Ramanan V, Kumaresan N. Comparative Study of Mental Stress Score between Various Surgical Modalities for Treatment of 10–20 mm Renal Calculi. *Int J Sci Stud* 2018;5(11):170-173.

Source of Support: Nil, **Conflict of Interest:** None declared.