

Neonates Born to Mothers with SARS-CoV-2 Infection: A Prospective Observational Study

Shweta Pathak¹, Asha Tiwari¹, Monica Lazarus²

¹Assistant professor, Department of Paediatrics, NSCB Medical College, Jabalpur, Madhya Pradesh, India, ²Associate professor, Department of Paediatrics, NSCB Medical College, Jabalpur, Madhya Pradesh, India

Abstract

Objectives: To assess the clinical presentation and risk of mother-to-infant transmission of SARS-CoV2 in infants born to mothers with COVID-19 infection.

Design: Prospective observational study.

Setting: Tertiary care centre and a dedicated covid hospital in central India.

Participants: Infants born to mothers diagnosed to have COVID-19 at the time of delivery, born between 1 June and 25 September 2020.

Results: 40 infants were born to mothers with COVID-19 at the time of delivery. Three neonates tested positive for SARS-CoV2 after birth-one had mild respiratory symptoms and the other two infants remained well. A preterm baby who was negative for covid died due to extreme prematurity with sepsis (early onset) with respiratory failure on day 4 of life and another baby who was negative died due to Hypoxic ischaemic encephalopathy stage 3 with Meconium aspiration syndrome with respiratory failure. Remaining 38 neonates and their mother roomed in while in hospital and all were breast fed. None of the other neonates developed any significant health issues or developed symptoms attributable to SARS CoV2.

Conclusions: The risk of mother-to-infant transmission of SARS-CoV2, in the perinatal period is very low. Breast feeding and rooming in can be practiced safely with adequate infection control precautions with negligible clinical risk to the infant.

Key words: Breastfeeding, COVID19, Newborn

INTRODUCTION

The novel SARS-CoV-2 which leads to the clinical syndrome now labeled as COVID-19 was first detected in the Wuhan – Hubei Province of China in December 2019. Since then, it has spread across the world with the WHO declaring it as a global pandemic on March 11, 2020.^[1] Knowledge about the epidemiology and clinical presentation of COVID-19 is rapidly evolving. Although the virus affects individuals across the age spectrum, it is becoming increasingly apparent that outcomes in adults are worse than in children.^[2,3] The vast

majority of subjects infected display only mild symptoms or remain asymptomatic.^[4] Women are deemed to be in an immunocompromised state and data from previous coronavirus epidemics (SARS-CoV and MERS-CoV) have shown that they were at a high risk of morbidity and mortality.^[5] The literature published so far on COVID-19, however, suggests that hospitalized pregnant women do not seem to be at a higher risk of adverse outcomes compared with hospitalized non-pregnant individuals.^[6,7] The risk of vertical transmission to infants born to mothers with COVID-19 seems low.^[7-11] In the few infants who tested positive following birth, it was not certain whether the transmission was vertical or postnatal and further, the majority of these infants had only mild-to-moderate disease.

Here, we present the short-term outcomes of infants born to mothers infected with SARS-CoV-2 and the safety of a policy of rooming in and breastfeeding such babies at our hospital.

Access this article online



www.ijss-sn.com

Month of Submission : 08-2020
Month of Peer Review : 09-2020
Month of Acceptance : 09-2020
Month of Publishing : 10-2020

Corresponding Author: Shweta Pathak, Department of Paediatrics, NSCB Medical College, Jabalpur, Madhya Pradesh, India.

MATERIALS AND METHODS

All neonates born to mothers who tested positive for SARS-CoV-2 during pregnancy from June 1, 2020, to September 20, 2020, were eligible to be included in the study. SARS-CoV-2 virus was diagnosed by real-time reverse transcriptase PCR in nasopharyngeal swab taken up by an expert on day 2 of life or at admission for neonates referred from other centers as COVID suspect. Interpretation of the result was performed according to the manufacturer's recommendation. Once a negative result was obtained, babies were labeled as COVID negative and test was not repeated for asymptomatic neonates as per the institutional protocols. Neonates who were well enough to be cared on the postnatal ward were allowed to room in with their mothers except where mother or child was too critical and required admission in intensive care unit. Neonates who roomed in were allowed to breastfeed with adequate droplet and contact precautions in accordance with recommendations from professional bodies. This included caring for the infant in a separate baby cot which is placed at least 6 ft from the mother's bed, mother wearing a mask when in close proximity to the infant and performing hand hygiene with either soap and water or alcohol-based gel prior to handling the baby. No visitors apart from health personnel with appropriate personal protective equipment were permitted into the rooms while in hospital. The parents were advised to continue the same precautions at home after discharge till the mother was considered negative for COVID-19 and had been in isolation for 14 days.

Data Collection and Analyses

The baseline characteristics (age at admission, sex, birth weight, and gestational age), clinical details which included term/preterm, clinical diagnosis (transient tachypnea of newborn), hypoxic ischemic encephalopathy Stage 3, meconium aspiration syndrome, hypoglycemia, sepsis (early onset), roomed in with mother in hospital, breastfed at the time of discharge, and results of COVID-19 screening were collected from the patient record. Informed consent was obtained from the parents for inclusion in the study. Data were collected on an electronic database and analyzed.

Table 1: Baseline characteristics

Baseline characteristics	Median (range)* Number (percentage)†
Day of admission	4 days (1 PND-22 days)
Birth weight in grams	2508 (1200–3600)*
Gestation (weeks)	38 weeks (28–42)*
Normal vaginal delivery	18 (45%)
Cesarean section	22 (55%)
Male:female	1.2:1 (22 M 18F)
Full term (AFD)	36 (90%)
Full term (SFD)	3 (7.5%)†
Preterm (SFD)	1 (2.5%)†

RESULTS

A total of 40 neonates born to COVID-positive mothers were included in the study period of June 1–September 25. The baseline characteristics are summarized in Table 1.

Out of the 40 neonates as summarized in Table 2, 7 were admitted to intensive care unit – most of these 5 (12.5%) for transient tachypnea and/or prematurity (2.5%). There were 2 (5%) deaths out of total 40 neonates born to COVID-positive mother till 25 of September both were negative for COVID (A preterm baby died due to extreme prematurity with sepsis (early onset) with respiratory failure on day 4 of life and another baby died due to hypoxic ischemic encephalopathy Stage 3 with meconium aspiration syndrome with respiratory failure. Thirty-eight of the 35 infants (95%) received breast milk either directly if rooming in with mother or through expressed breast milk if admitted to NICU. At the time of writing the report, the discharge outcomes of 24 neonates were known while 8 infants are still in the hospital.

Three neonates tested positive [Table 3] for SARS-CoV-2, out of these 3 the first one who was born through LSCS admitted for mid cough and fever on day 20 of life and a history of contact with COVID-positive mother after almost 15 days of her delivery came positive for COVID-19, received supportive treatment, and discharged successfully on day 10 of isolation in stable condition, and other two came positive for COVID-19 on day 3 and day 10 of life both were asymptomatic discharged successfully after 10 days of isolation.

Table 2: Indication for admission in HDU

Indication for admission in HDU (isolation ward)	Number (percentage)
Indication for admission in HDU	
Prematurity	1 (2.5)†
Transient tachypnea of newborn	5 (12.5)†
Hypoxic ischemic encephalopathy stage 3	1 (2.5)†
Meconium aspiration syndrome	1 (2.5)
Hypoglycemia	2 (5)†
Sepsis (early onset)	2 (5)

†Number (percentage)

Table 3: Short-term outcome of neonates

Neonates positive for RT-PCR	3 (7.5)
Roomed in and breastfed with mother†	38 (95)
Discharged	28 (70)
Still admitted	10 (25)
Mortality	2 (5)
HIE stage 3 meconium aspiration syndrome	1 (2.5)
Extreme prematurity with NEC Stage 3 with septic shock	1 (2.5)

DISCUSSION

Despite initial concern that pregnant women and the newborn may be high-risk groups compared with the general population based on outbreaks of other coronavirus diseases in the past, it has become increasingly clear that this is not the case with the SARS-CoV-2 pandemic.^[6-11] The risk of vertical transmission of the virus from mother to infant before or during delivery has been shown to be low.^[7-12] This has been confirmed based on virus testing and clinical features in our cohort, with the estimated risk being only 7.5% (3/35). This figure seems to be similar to previously published data.^[6,9,13] The low risk of vertical transmission has been hypothesized to be due to paucity of ACE2 receptors in the placenta which may be necessary for transplacental transfer to the fetus.^[9] Regardless of the above, it would appear from our series that the short-term clinical risks to the infant from maternal COVID-19 at the time of delivery are minimal. Due to the uncertainty surrounding the outcomes of mothers affected by COVID-19 and their infants, recommendations on the postnatal management of the mother-infant dyad from professional bodies have been inconsistent.^[12,14-17] Some guidelines advocate caring for the affected mothers and their infants in separate rooms when feasible to reduce the risk of mother-infant transmission postnatally and also recommend avoiding direct breastfeeding while mother is still infected, unless mother expresses her wish to directly breastfeed.^[16,17] At our hospital, we found that the benefits of both rooming in and breastfeeding with good infection control precautions, for mother-infant bonding and long-term breastfeeding far outweighed the small risk of mother-infant transmission. Thus, we strongly recommended rooming in and direct breastfeeding for all well mother-infant dyads while in hospital as well as after discharge. Our study has validated this approach with no clinically or laboratory-proven mother to infant transmission of the virus during the hospital stay, even with a very high rate of breastfeeding in the discharged infants.

To the best of our knowledge, this study is the single largest series on the outcomes of infants born to mothers with COVID-19 with follow-up of their health status post-discharge. However, our study has limitations. The diagnosis of COVID-19 and virus carriage was based on NP swabs. Although the absolute sensitivity of detecting SARS-CoV2 with NP swabs is unknown, the modality is only around 70% sensitive for diagnosing respiratory viral infections.^[18] The universally good clinical outcomes in all the infants are nevertheless encouraging. The follow-up of infants was conducted by telephonic interview of the parents. There is a chance that asymptomatic and mildly symptomatic infants infected with the virus may have been missed in the absence of testing, although we would suggest that this is of limited clinical significance.

CONCLUSION

The risk of mother-to-infant transmission of SARS-CoV-2, in the perinatal period, is very low. Breastfeeding and rooming in can be practiced safely with adequate infection control precautions. The risk of adverse outcome to infants born to mothers who have SARS-CoV-2 infection at birth is minimal aside from the risk of premature delivery due to iatrogenic/maternal causes. However, there are no published long-term outcome data on these infants and further follow-up studies will be needed to fully ascertain adverse outcomes in this group of infants.

ACKNOWLEDGMENT

We would like to acknowledge our respected Head of the Department, Dr. Avyact Agrawal sir and Dr. Pawan sir for their guidance. Our colleagues Dr. Pratibha Bamne, Dr. Ravi Uikey, and Dr. Akhilendra Parihar, Senior Residents and Postgraduate residents for their constant efforts in the management of COVID children. Special thanks to Dr. Shashank Pandey for literature search.

REFERENCES

1. World Health Organization. Corona Virus Disease 2019 (COVID-19): Situation Report 51. Geneva: World Health Organization; 2003. Available from: https://www.who.int/docs/default-source/coronaviruse/situation-reports/20200311-sitrep-51-covid19.pdf?sfvrsn=1ba62e57_10. [Last accessed on 2020 Jul 11].
2. Zhou F, Yu T, Du R, Fan G, Liu Y, Liu Z, *et al.* Clinical course and risk factors for mortality of adult inpatients with COVID-19 in Wuhan, China: A retrospective cohort study. *Lancet* 2020;395:1054-62.
3. Ludvigsson JF. Systematic review of COVID-19 in children shows milder cases and a better prognosis than adults. *Acta Paediatr* 2020;109:1088-95.
4. Long L, Zeng X, Zhang X, Xiao W, Guo E, Zhan W, *et al.* Short-term outcomes of COVID-19 and risk factors for progression. *Eur Respir J* 2020;55:2000990.
5. Guan WJ, Liang WH, Zhao Y, Liang HR, Chen ZS, Li YM, *et al.* Comorbidity and its impact on 1590 patients with COVID-19 in China: A nationwide analysis. *Eur Respir J* 2020;55:2000547.
6. Zaigham M, Andersson O. Maternal and perinatal outcomes with COVID-19: A systematic review of 108 pregnancies. *Acta Obstet Gynecol Scand* 2020;99:823-9.
7. Huntley BJ, Huntley ES, Di Mascio D, Chen T, Berghella V, Chauhan SP. Rates of maternal and perinatal mortality and vertical transmission in pregnancies complicated by severe acute respiratory syndrome coronavirus2 (SARS-CoV-2) infection: A systematic review. *Obstet Gynecol* 2020;136:303-12.
8. Akhtar H, Patel C, Abuelgasim E, Harky A. COVID-19 (SARS-CoV-2) infection in pregnancy: A systematic review. *Gynecol Obstet Invest* 2020;2020:1-12.
9. Eglhoff C, Vauloup-Fellous C, Picone O, Mandelbrot L, Roques P. Evidence and possible mechanisms of rare maternal-fetal transmission of SARS-CoV-2. *J Clin Virol* 2020;128:104447.
10. Chen H, Guo J, Wang C, Luo F, Yu X, Zhang W, *et al.* Clinical characteristics and intrauterine vertical transmission potential of COVID-19 infection in nine pregnant women: A retrospective review of medical records. *Lancet* 2020;395:809-15.
11. Schwartz DA. An analysis of 38 pregnant women with COVID-19, their newborn infants, and maternal-fetal transmission of SARS-CoV-2:

- Maternal coronavirus infections and pregnancy outcomes. Arch Pathol Lab Med 2020;2020:5858.
12. Royal College of Paediatrics and child Health. COVID-19 Guidance for Neonatal Settings; 2020.
 13. Juan J, Gil MM, Rong Z, Zhang Y, Yang H, Poon LC. Effect of coronavirus disease 2019 (COVID-19) on maternal, perinatal and neonatal outcome: Systematic review. Ultrasound Obstet Gynecol 2020;56:15-27.
 14. Davanzo R, Moro G, Sandri F, Agosti M, Moretti C, Mosca F. Breastfeeding and coronavirus disease-2019: Ad interim indications of the Italian society of neonatology endorsed by the union of European neonatal and perinatal societies. Matern Child Nutr 2020;16:e13010.
 15. Centre for Disease Control and Prevention. Evaluation and Management Considerations for Neonates at Risk for COVID-19; 2020. Available from: <https://www.cdc.gov/coronavirus/2019-ncov/hcp/caring-for-newborns.html>. [Last accessed on 2020 Jul 11].
 16. American Academy of Pediatrics. Management of Infants Born to Mothers with Suspected or Confirmed COVID-19; 2020. Available from: <https://www.services.aap.org/en/pages/2019-novel-coronavirus-covid-19-infections/clinical-guidance/faqs-management-of-infants-born-to-covid-19-mothers>. [Last accessed on 2020 Jul 11].
 17. Wang L, Shi Y, Xiao T, Fu J, Feng X, Mu D, *et al*. Chinese expert consensus on the perinatal and neonatal management for the prevention and control of the 2019 novel coronavirus infection (First edition). Ann Transl Med 2020;8:47.
 18. Lieberman D, Lieberman D, Shimoni A, *et al*. Identification of respiratory viruses in adults: nasopharyngeal versus oropharyngeal sampling. J Clin Microbiol 2009;47:3439-43.

How to cite this article: Pathak S, Tiwari A, Lazarus M. Neonates Born to Mothers with SARS-CoV-2 Infection: A Prospective Observational Study. Int J Sci Stud 2020;8(7):69-72.

Source of Support: Nil, **Conflicts of Interest:** None declared.