

# Clinical Profile of Granuloma Brain: A Prospective Study

A Thomas Edwin Raj<sup>1</sup>, P K Murugan<sup>2</sup>, S Saravanan<sup>2</sup>, M Radha<sup>3</sup>, Nancy Dora<sup>4</sup>, Heber Anandan<sup>5</sup>

<sup>1</sup>Senior Resident, Department of Neurology, Tirunelveli Medical College and Hospital, Tirunelveli, Tamil Nadu, India, <sup>2</sup>Professor, Department of Neurology, Tirunelveli Medical College and Hospital, Tirunelveli, Tamil Nadu, India, <sup>3</sup>Assistant Professor, Department of Neurology, Tirunelveli Medical College and Hospital, Tirunelveli, Tamil Nadu, India, <sup>4</sup>Professor, Department of Radiology, Tirunelveli Medical College and Hospital, Tirunelveli, Tamil Nadu, India, <sup>5</sup>Senior Clinical Scientist, Department of Clinical Research, Dr. Agarwal's Health Care Limited, Tirunelveli, Tamil Nadu, India

## Abstract

**Introduction:** Granuloma of the central nerve system is the most common radiological abnormality seen in patients with acute onset seizures in India. Neurocysticercosis (NCC) and tuberculoma are the most common causes of this granuloma.

**Aim:** To study the clinical feature of granuloma brain and the effect of treatment on follow-up.

**Materials and Methods:** Patients presenting with radiological evidence of single ring enhancing lesions are taken up for study. Physiological calcifications, calcified granulomas, patients with primary malignancy, and HIV were excluded.

**Results:** Granuloma is more common in the age group from 11 to 30 years with female predominance. Focal seizures are common and parietal lobe is the most common location. Perilesional edema is more common in tuberculoma. Tuberculoma has a thick and irregular wall. Patients with NCC were treated with albendazole, steroid, and antiepileptic drugs (AEDs), patients with tuberculoma were treated with antituberculous therapy, steroids, and AEDs. Follow-up imaging at 6 and 12 months showed complete resolution of cysticercosis granuloma and tuberculoma in most of the patients.

**Conclusion:** Focal seizures are the most common seizure type followed by generalized seizures. Majority of the granuloma occurred in parietal lobes. Follow-up computed tomography at 6 and 12 months revealed complete resolution of lesions in most of NCC patients and resolving lesions in most of the tuberculoma patients.

**Key words:** Antiepileptic drugs, Antituberculous therapy, Neurocysticercosis, Tuberculoma

## INTRODUCTION

Granuloma of the central nervous system is the most common radiological abnormality seen in patients with acute-onset seizures in India and many other developing countries. Neurocysticercosis (NCC) is the most likely cause of this granuloma. A single degenerating cyst<sup>1</sup> is the most frequent finding associated with NCC in the Indian subcontinent. Single cysticercus granuloma is one which measures <20 mm in diameter, may be associated with cerebral edema not severe enough to produce midline

shift, and occur in patients with seizures and normal neurological status, without evidence of active systemic disease. When this granuloma resolves spontaneously, either disappearing or changing into a calcified nodule, the diagnosis of NCC is very likely. The second most common cause of these computed tomography (CT)-detected granuloma is tuberculoma, in patients with these granuloma similar clinical and neuroimaging features are also present. Granuloma visualized on CT scanning is the most common radiological abnormality in Indian patients with new-onset seizures.<sup>2</sup> In 1980, Tandon and Bhargava<sup>3</sup> first reported these lesions; at that time these CT-enhancing lesions were presumed to be tuberculoma and often were treated with empirical anti-tuberculous drugs. Subsequently, histopathological studies of brain tissue biopsy samples have suggested that, in majority, CT-enhancing lesions represent dying<sup>4</sup> cysticercal lesions (larval stage of tapeworm *Taenia solium*).

### Access this article online



www.ijss-sn.com

**Month of Submission :** 03-2017  
**Month of Peer Review :** 04-2017  
**Month of Acceptance :** 05-2017  
**Month of Publishing :** 05-2017

**Corresponding Author:** Dr. P K Murugan, Department of Neurology, Tirunelveli Medical College and Hospital, Tirunelveli, Tamil Nadu, India. Phone: +91-9486306399. E-mail: thomasedwinraj2015@gmail.com

In a developing country like ours where both tuberculosis and cysticercosis are common, it is difficult to differentiate between tuberculoma and a cysticercal granuloma. Some granuloma “heal” by becoming calcified. These patients require only antiepileptic therapy, and this medication may be withdrawn safely after the lesion has resolved on CT scanning. Anticysticercal drugs given for the enhancing lesions for 14 days with steroids and antituberculous therapy (ATT) is administered for the tuberculous granulomas. For patients presenting with a headache and other symptoms, symptomatic therapy was instituted.

### Aim

To study the clinical feature of granuloma brain and the effect of treatment on follow-up.

## MATERIALS AND METHODS

This prospective observational study was conducted in Department of Neurology at Tirunelveli Medical College and Hospital. All patients presenting with radiological evidence of single ring enhancing granuloma of brain were included in the study.

### Exclusion Criteria

Patients with physiological calcification and calcified granulomas, neonates and infants, patients who are pregnant, patients with claustrophobia, patients with primary malignancy, and patients who are HIV positive. The study was commenced after obtaining approval from the Institutional Ethical Committee. Written informed consent was obtained from those who were willing to participate in the study after explaining to them about the nature of the study, additional sequences of imaging and the need for follow-up in the prescribed format and in the regional language. Left thumb impression was obtained from those patients who are illiterate. A detailed history was obtained from patients presenting with imaging evidence of single ring enhancing granuloma brain. All patients were treated with antiepileptic drugs (AEDs) for the entire study period. A course of steroids of 1 mg/kg was started on patients who had signs of increased intracranial tension or imaging evidence of profound perilesional edema. No serological or cerebrospinal fluid tests for NCC were included in the study. All patients were followed up during the entire study period. They were asked to report for recurrence of seizures in addition to a monthly follow-up. CT scans with contrast were done at the end of 6 and 12 months for regression, persistence, enlargement, resolution, and calcification.

## RESULTS

A total of 42 patients were included in the study; granuloma is more common in the age group from 11 to 20 years. Second most common age group is from 21 to 30 years. Granuloma is more common in the female is probably due to gender bias in seeking medical attention (Table 1).

In this study, most of the patients presented with focal seizures followed by seizures of generalized from the onset (Table 2). Granuloma is predominantly in the parietal region. The second most common region is over the frontal region (Table 3). Common size of granuloma is between 10 and 20 mm. The second most common is <10 mm (Table 4).

In this study, perilesional edema was seen in 20 patients. Thick and irregular wall sign of tuberculoma was seen in 18 patients (Figure 1). Thin and regular wall sign of cysticercosis was seen in 24 patients (Figure 2).

**Table 1: Age distribution in study patients**

Age	Patients
<10	5
11-20	13
21-30	10
31-40	8
41-50	3
>50	3

**Table 2: Seizure semiology at presentation**

Types of seizure	Patients
Focal seizures	24
Complex partial seizures	2
Generalized from onset	16

**Table 3: Location of the granuloma**

Location	Patients
Frontal	11
Parietal	20
Temporal	3
Occipital	4
Cerebellum	2
Brain stem	2

**Table 4: Size of the granuloma**

Size (mm)	Patients
<10	14
10-20	16
>20	12




Figure 1: Magnetic resonance imaging brain showing lesion with thick irregular margin




Figure 3: Magnetic resonance imaging brain with contrast showing conglomerate ring enhancing lesion




Figure 2: Magnetic resonance imaging brain with contrast showing small ring enhancing lesion with smooth wall over right parietal lobe




Figure 4: Computed tomography brain showing left parietal granuloma with perilesional edema


A conglomerate lesion indicates tuberculoma (Figure 3). A calcified lesion with edema was seen in patients (Figure 4). The presence of scolex in the lesions indicates cysticercosis granuloma (Figure 5).

On follow-up, the disappearance of tuberculosis granuloma was noted after 9 months in 14 patients, 2 patients required 18-24 months course, 2 patients required prolonged treatment course for more than 2 years. After 6 months follow-up CT showed the disappearance of cysticercosis granuloma in 24 patients.

## DISCUSSION

A total of 42 patients with single ring enhancing lesions were taken up for study. Single enhancing lesions according to Rajshekhar and Chandy is predominately affects males the age group of 20-30 years.<sup>5</sup> In our study, it is 11-30 years

and predominately female population. The greater incidence in our region is due to gender bias in seeking medical attention. Focal seizure is most common seizure type in our study which is in accordance with Chopra *et al.* study.<sup>6</sup> In our study granuloma predominately affects the parietal regions and frontal regions. Brain stem and cerebellum involvement were observed in 2 patients each. 24 patients with single ring enhancing granulomas were due to NCC and 18 were due to tuberculoma. Patients are having lesion more than 20 mm in size are taken as definite tuberculomas, <10 mm in size are taken as NCC and size between 10 and 20 mm with irregular wall and severe perilesional edema are taken as tuberculomas. Conglomerate lesions are common in tuberculoma patients. Single granuloma measuring <20 mm in diameter, regular outline, may be associated with perilesional edema not severe enough to produce midline shift, is taken as cysticercal granuloma. Tuberculoma by contrast is usually irregularly shaped solid and more than 20 mm in



**Figure 5: Axial fluid-attenuated inversion recovery image showing scolex**

size. They are often associated with severe perilesional edema, midline shift, and focal neurological deficit. Fluid-attenuated inversion recovery suppression of signal core in ring enhancing lesion rules out tuberculoma and seen in NCC.

There is a wide variation in the rates of complete resolution mentioned by different authors<sup>7,8</sup> ranging from 22% to 100% at 6 months/12 months after the first CT scan. In this study, the follow-up contrast CT revealed a complete resolution in 18 patients (75%) at 6 months and 21 patients (88%) at 1 year in NCC patients. Patients treated with ATT for tuberculoma in the partial resolution were present in 12 patients (66%) and complete resolution in 16 (88%) patients at 1 year. For patients without resolution at 1 year duration of ATT was extended for 15-18 months. An extended period of ATT was given more than 2 years for 2 patients. AEDs were given for 2-3 years for tuberculoma. AEDs are given for patients with cysticercus granuloma till the disappearance of lesions.<sup>9</sup> Injection dexamethasone 8mg twice daily given during the acute symptoms and a tapering dose of prednisolone 1 mg/kg for 8 weeks for tuberculoma and 2 weeks for cysticercosis. Symptomatic

treatment was given for a headache and other non-specific complaints.<sup>10</sup>

## CONCLUSION

Focal seizures are the most common seizure type followed by generalized seizures. Majority of the granuloma occurred in parietal lobes. Follow-up CT at 6 and 12 months revealed complete resolution of lesions in most of NCC patients and resolving lesions in most of the tuberculoma patients. Patients treated with injection streptomycin are associated with greater resolution of lesions. An extended period of ATT more than 2 years may be required for resistance tuberculoma. AEDs are required till complete resolution of lesions in NCC and for 2-3 years in tuberculous granulomas.

## REFERENCES

1. Rajshekhar V. Etiology and management of single small CT lesions in patients with seizures: Understanding a controversy. *Acta Neurol Scand* 1991;84:465-70.
2. Kumar Garg R, Kumar Singh M, Misra S. Single-enhancing CT lesions in Indian patients with seizures: A review. *Epilepsy Res* 2000;38:91-104.
3. Tandon PN, Bhargava S. CNS tuberculosis lessons learnt from CT studies. *Neurol India* 1980;28:225-30.
4. Rajshekhar V, Oommen A. Serological studies using ELISA and EITB in patients with solitary cysticercus granuloma and seizures. *Neurol Infect Epidemiol* 1997;2:177-80.
5. Rajshekhar V, Chandy MJ, editors. Clinical manifestation of solitary cysticercus granuloma. *Cysticercus Granuloma: The Disappearing Lesion*. Hyderabad: Orient Longman; 2000. p. 29-39.
6. Chopra JS, Sawhney IM, Suresh N, Prabhakar S, Dhand UK, Suri S. Vanishing CT lesions in epilepsy. *J Neurol Sci* 1992;107:40-9.
7. Del Brutto OH, Roos KL, Coffey CS, Garcia HH. Meta-analysis: Cysticidal drugs for neurocysticercosis: Albendazole and praziquantel. *Ann Intern Med* 2006;145:43-51.
8. Sethi PK, Kumar BR, Madan VS, Mohan V. Appearing and disappearing CT scan abnormalities and seizures. *J Neurol Neurosurg Psychiatry* 1985;48:866-9.
9. Mukherjee A, Chakravarty A. Partial and complex partial seizures. In: Singhal BS, Nag D, editors. *Epilepsy in India 2000*. Mumbai: Indian Epilepsy Association; 2000. p. 50-75.
10. Garg RK, Misra S. Non-epileptic manifestations in patients with single enhancing computed tomography lesions. *J Assoc Physicians India* 2000;48:788-93.

**How to cite this article:** Raj ATE, Murugan PK, Saravanan S, Radha M, Dora N, Anandan H. Clinical Profile of Granuloma Brain: A Prospective Study. *Int J Sci Stud* 2017;5(2):1-4.

**Source of Support:** Nil, **Conflict of Interest:** None declared.